

# LIV-PRO<sup>™</sup>

## *Dosage Directions*

MLF Biotech, Inc.  
P.O. Box 538  
Milford, IN 46542-0538

Phone: (574) 658-2266  
Fax: (574) 658-2246  
www.mlfbiochem.com

**For best results, Liv-Pro<sup>™</sup> should be administered as follows:**

We recommend administering Liv-Pro<sup>™</sup> through drinking water so that the live cells may be homogenously dispersed throughout. If feed is used, the distribution of the beneficial bacteria is questionable. Additionally, sick birds do not eat much feed, but will still drink water.

*When chlorinating water, do not use product. Drinking water must be changed every 24 hours.*

**Broilers and Ducks:**

**1 to 14 days old** - 1 gram/1000 birds/day

Dissolve 50 grams into 384 gallons of drinking water (non-chlorinated).

**Over 14 days old** - 2 grams/1000 birds/day

Dissolve 100 grams into 384 gallons of drinking water (non-chlorinated).

**Layers and Breeders:**

**3 - 5 grams per 1,000 birds/day** (depending on health status of flock)

For a healthy flock, dissolve 150 grams into 384 gallons of drinking water (non-chlorinated).

For an unhealthy flock, dissolve 250 grams into 384 gallons of drinking water (non-chlorinated).

**Turkeys:**

**5 grams per 1,000 birds/day**

Dissolve 250 grams into 384 gallons of drinking water (non-chlorinated).

**Metering systems:**

At a rate of 1 gram of Liv-Pro<sup>™</sup> per 1,000 birds, add 50 grams of Liv-Pro<sup>™</sup> to 5 gallons of water (use dosage rates above). Dissolve Liv-Pro<sup>™</sup> in 2 gallons or less of solution to be metered at a ratio that ensures it gets it used up in 4 hours (to avoid any contamination and/or efficacy loss). Set proportioner to meter one ounce per gallon of water.